
integral	ass 540 is considered to be an l part of Class 260 (see the Class edule for the position of this n schedule hierarchy). This Class	12	Plural oxygens in both rings which share a spiro atom (e.g., 17,20;20,21 bismethylenedioxy-pregnanes, etc.)
retains	all pertinent definitions and ines of Class 260.	13	Nitrogen, sulfur, cyano or -C(=X)-, wherein X is chalcogen, bonded directly to the cyclopentanohydrophenanthrene ring system
	ORGANIC COMPOUNDS (CLASS 532,	14	Halogen attached directly
1	SUBCLASS 1) .HETEROCYCLIC CARBON COMPOUNDS CONTAINING A HETERO RING HAVING CHALCOGEN (I.E., OXYGEN, SULFUR, SELENIUM OR		or indirectly to the cyclopentanohydrophenanthrene ring system by acyclic nonionic bonding
	TELLURIUM) OR NITROGEN AS THE ONLY RING HETERO ATOMS	15	The cyclopentanohydrophenanthrene
2	Cyclopentanohydrophenanthrene ring system containing		ring system is part of a polycyclo ring system having at least five cyclos
3	Heavy metal or aluminum containing	16	Hetero ring is one of the cyclos of the polycyclo ring
4	Boron or silicon containing		system
5	Phosphorus attached directly or indirectly to the cyclopentanohydrophenanthrene ring system by nonionic bonding	17	The hetero ring is five- membered, consisting of one oxygen and four carbons, and shares the spiro atom with a six-membered oxygen containing
6	Spiro		hetero ring (e.g., sapogenins,
7	Plural spiro atoms		etc.)
8	The	18	Purification or recovery
9	cyclopentanohydrophenanthrene ring system is part of a polycyclo ring system having at least five cyclosNitrogen containing hetero	19	Chalcogen bonded directly at the 12-position of the cyclopentanohydrophenanthrene ring system (e.g., hecogenin, etc.)
	ring as one of the cyclos of	20	Chalcogen bonded directly
10	the polycyclo ring systemThe cyclopentanohydrophenanthrene		at the 11-position of the cyclopentanohydrophenanthrene ring system
	ring system shares spiro atoms with two hetero rings, each of which contains two oxygens (e.g., 3,17-bis-ketals, etc.)	21	Chalcogen bonded directly at the 7-position of the cyclopentanohydrophenanthrene ring system
11	The	22	
	cyclopentanohydrophenanthrene ring system shares a spiro atom with a lactone ring (i.e., -C(=X)-O- is part of	22	or sulfur bonded directly to the cyclopentanohydrophenanthrene ring system
	the ring, wherein X is chalcogen)	23	The spiro atom is the 17- position carbon of the cyclopentanohydrophenanthrene ring system



53		69	Halogen bonded directly to the
	directly at the 17-position of the		cyclopentanohydrophenanthrene ring system
	cyclopentanohydrophenanthrene ring system	70	0xygen bonded directly at the 11-position of the
54	The hetero ring consists of two nitrogens and three		cyclopentanohydrophenanthrene ring system
	carbons and is ortho fused to the D ring	71	0xygen attached directly to the B ring or indirectly to
55	Chalcogen in the hetero ring		the A or B ring by acyclic nonionic bonding
56	<pre>The hetero ring is five- membered</pre>	72	The hetero ring is a lactone (i.e., containing $-C(=X)-0-$ as
57	The hetero ring is ortho- fused to the A ring		<pre>part of the ring, wherein X is chalcogen)</pre>
58	<pre>The hetero ring is five- membered</pre>	73	The lactone ring shares at least three ring members with
59	The hetero ring contains sulfur		one other cyclo of the polycylo ring system (i.e.,
60	The hetero ring is a cyclic		bridged)
	anhydride (i.e., containing -	74	The lactone ring shares
	C(=X)-O-C(=Y)- as part of the		the 11,12,13-positions of the
	ring, wherein X and Y are		cyclopentanohydrophenanthrene
	chalcogen; e.g., 5,8-maleic		ring system (e.g., 11, 18-
	anhydride adduct of 5,7,9(11)-		lactones, etc.)
	<pre>pregnatrien-3,20-di-one, etc.)</pre>	75	The lactone ring shares a
61	The hetero ring contains		ring carbon with two other
	plural oxygens		cyclos of the polycyclo ring
62	Two of the cyclos share at		system (e.g., peri-fused,
	least three ring members or a	7.6	etc.)
	ring carbon is shared by three	76	The hetero ring is three-
	of the cyclos (e.g., bridged,		membered consisting of one
	peri-fused, etc.)		oxygen and two carbons (e.g.,
63	The hetero ring is ortho-	77	oxirane, etc.)
- 1	fused to the D ring	1 1	The polycyclo ring system contains plural oxirane rings
64	At least six cyclos in the	78	
	polycyclo ring system	70	The hetero ring shares the
65	Nitrogen or acyclic		1,2-positions of the cyclopentanohydrophenanthrene
	chalcogen bonded directly to		ring system
	the hetero ring (e.g., cyclic	79	The hetero ring shares the
	carbonates, etc.)	13	4,5-positions of the
66	The A ring is a benzene ring		cyclopentanohydrophenanthrene
67	Sulfur or nitrogen	80	ring system
	attached directly or	80	The hetero ring shares the
	indirectly to the		5,6-positions of the
	cyclopentanohydrophenanthrene		cyclopentanohydrophenanthrene
	ring system by acyclic	0.1	ring system
	nonionic bonding	81	The hetero ring shares the
68	Halogen attached		6,7-positions of the
	indirectly to the		cyclopentanohydrophenanthrene
	cyclopentanohydrophenanthrene		ring system
	ring system by acyclic		
	nonionic bonding		

82	The hetero ring shares the 11,12-positions of the cyclopentanohydrophenanthrene	94	Hetero ring attached directly to the cyclopentanohydrophenanthrene
83	ring systemThe hetero ring shares the 14,15-positions of the cyclopentanohydrophenanthrene ring system	95 96	ring system by nonionic bondingThe hetero ring contains nitrogenPlural nitrogen containing
84	The hetero ring shares the 16,17-positions of the cyclopentanohydrophenanthrene ring system	90	hetero rings bonded directly to the cyclopentanohydrophenanthrene ring system
85	Halogen bonded directly to the cyclopentanohydrophenanthrene ring system	97	The hetero ring is bonded directly at the 3-position of the cyclopentanohydrophenanthrene
86 87	Saturated A ringThe hetero ring shares the 9,11-positions of the	98	ring systemThe A ring is a benzene ring
88	cyclopentanohydrophenanthrene ring systemHalogen bonded directly to	99	Halogen bonded directly to the cyclopentanohydrophenanthrene
	the cyclopentanohydrophenanthrene ring system	100	ring systemThe hetero ring contains plural chalcogens
89	is chalcogen, bonded directly to the cyclopentanohydrophenanthrene ring system	101	The hetero ring and acyclic chalcogen are both bonded directly at the 17 position of the
90	The hetero ring shares at least three ring members with one other cyclo of the polycyclo ring system (i.e., bridged)	102	<pre>cyclopentanohydrophenanthrene ring systemThe hetero ring contains - C(=X)-O-, wherein X is chalcogen, as part of the ring (e.g., lactones, etc.)</pre>
91	Bridge consisting of oxygen and carbon between the 6- and 10-positions of the cyclopentanohydrophenanthrene	103	<pre>Additional chalcogen, cyano, or -C(=X)-, wherein X is chalcogen, bonded directly to the hetero ring</pre>
92	ring system (e.g., 6,19-oxido steroids, etc.)Bridge consisting of oxygen and carbon between the 11- and 13-positions of the	104	Nitrogen or sulfur attached directly or indirectly to the cyclopentanohydrophenanthrene ring system by nonionic bonding
	cyclopentanohydrophenanthrene ring system (e.g., 11,18-oxido steroids, etc.)	105	Chalcogen bonded directly at the 14-position of the cyclopentanohydrophenanthrene
93	The hetero ring shares a ring carbon with two other cyclos of the polycyclo ring		ring system or double bond in the D ring (e.g., cardenolides, etc.)
	system (e.g., peri-fused, etc.)	106	Nitrogen attached directly to the cyclopentanohydrophenanthrene ring system by nonionic bonding

107	Nitrogen containing hetero ring attached indirectly to the cyclopentanohydrophenanthrene ring system by nonionic	120	Chalcogen attached indirectly to the cyclopentanohydrophenanthrene ring system by nonionic bonding
	bonding	121	Azaporphyrins
108	The hetero ring is five-	122	Phthalocyanines
	membered and has plural hetero	123	Hetero ring attached directly
	atoms		or indirectly to the
109	The hetero ring is in the 17-		phthalocyanine ring system by
	position substituent of the		nonionic bonding
	cyclopentanohydrophenanthrene	124	The hetero ring is six-
	ring system		membered having nitrogen as a
110	The hetero ring is bonded		ring member
	directly to a $-C(=X)$ - group,	125	Plural hetero atoms in the
	wherein X is chalcogen		six-membered hetero ring
111	Having $-C(=X)-$, wherein X is	126	Triazines (including
	chalcogen, bonded directly at		hydrogenated)
	the 17-position of the	127	The hetero ring is five-
	cyclopentanohydrophenanthrene		membered having plural hetero
	ring system		atoms, at least one of which
112	Chalcogen or nitrogen in		is nitrogen
	chain between the hetero ring	128	Boron, germanium, phosphorus
	and the		or silicon containing
	cyclopentanohydrophenanthrene	129	Having $-C(=X)-$, wherein X is
	ring system		chalcogen, bonded directly to
113	Chalcogen in chain between		ring carbon of the
	the hetero ring and the		phthalocyanine ring system
	cyclopentanohydrophenanthrene		(e.g., tetracarboxy copper
	ring system		phthalocyanine, etc.)
114	Oxygen containing hetero ring	130	Having $-C(=X)-$, wherein X is
	attached indirectly to the		chalcogen, attached indirectly
	cyclopentanohydrophenanthrene		to ring carbon of the
	ring system by nonionic bonding		phthalocyanine ring system by
115	The hetero ring contains -		nonionic bonding (e.g.,
113	C(=X)-O-, wherein X is		<pre>phthalocyanine acetic acids, etc.)</pre>
	chalcogen, as part of the ring	131	,
	(e.g., lactones, etc.)	131	Sulfonyl bonded directly to ring carbon of the
116	Additional hetero atom in the		phthalocyanine ring system
	oxygen containing hetero ring	132	Chalcogen bonded directly to
117	The A ring is a benzene ring	132	the sulfonyl group
118	The hetero ring is bonded	133	Nitrogen bonded directly to
	directly to chalcogen which is	133	the sulfonyl group
	bonded directly to the	134	Additional nitrogen in the
	cyclopentanohydrophenanthrene	131	sulfonyl containing
	ring system		substituent
119	The chalcogen is bonded	135	Nitrogen attached indirectly
	directly at the 17-position of	133	to ring carbon of the
	the		phthalocyanine ring system by
	cyclopentanohydrophenanthrene		acyclic nonionic bonding
	ring system	136	Halogen bonded directly to
			ring carbon of the
			phthalocyanine ring system

137	At least eight halogens bonded directly to ring carbons of the phthalocyanine ring system	217	Double bond between the 2,3-positions of the bicyclo ring system (e.g., 2 cephem, etc.)
138	Processes of halogenating the phthalocyanine ring system	218	Ring expansion to produce the bicyclo ring system
139	Metal containing	219	7-amino cephalosporanic
140	Heavy metal or aluminum containing		<pre>acid per se or salt thereof (i.e., 7-ACA or salt thereof)</pre>
141	Specified crystalline form	220	Purification or recovery
	or processes of milling (e.g.,	221	7,7-disubstituted
	<pre>alpha crystalline form, ball milling, acid milling, etc.)</pre>	222	Additional hetero ring containing
142	Processes of forming the	223	\dots 2- or 4-position
	phthalocyanine ring system		substituent contains hetero
143	From reactant which		ring
	contains plural cyano groups	224	3-position substituent
	(e.g., preparing from		contains a pyridine ring
	<pre>phthalonitrile, etc.)</pre>		(e.g., quinoline,
144	From reactant which contains plural carbonyl		<pre>thienopyridine, lutidines, etc.)</pre>
	groups (e.g., preparing from	225	7-position substituent
	phthalic anhydride, etc.)		contains hetero ring
145	Porphyrins (including	226	3-position substituent
	hydrogenated; e.g.,		contains sulfur
	chlorophyll, etc.)	227	7-position substituent
200	Hetero ring is four-membered		contains hetero ring
	containing nitrogen and having	228	Alkyl, hydroxyalkyl,
	chalcogen double bonded		alkoxyalkyl or
	directly to a ring carbon which is adjacent to the ring		alkanoyloxyalkyl bonded
	nitrogen	0.00	directly to 3 position
201	Heavy metal containing	229	Sulfur containing
202	Plural hetero atoms in the	020	substituent
202	hetero ring	230	Alkyl, hydroxyalkyl,
203	Polycyclo ring system		alkoxyalkyl or
203	containing the hetero ring as		alkanoyloxyalkyl bonded directly to 3 position
	one of the cyclos	300	The six-membered ring
204	The ring nitrogen is shared	300	contains oxygen
201	by a ring containing at least	301	1-oxa-5-aza-
	seven members	301	bicyclo(4.2.0)octane
205	The ring nitrogen is shared		(including unsaturated)
	by a six-membered ring	302	The ring nitrogen is shared
214	The six-membered ring	302	by a five-membered ring
	contains sulfur	303	The five-membered ring
215	1-thia-5-aza-	303	contains an additional hetero
	bicyclo(4.2.0)octane		atom
	(including unsaturated; e.g.,	304	1-thia-4-aza-
	cepham, etc.)	-	bicyclo(3.2.0)hep-tane
216	The 1-thia-5-aza-		(including unsaturated; e.g.,
	bicyclo(4.2.0)oct-ane is part		penam, etc.)
	of a polycyclo ring system	305	The 1-thia-4-aza-
	having at least three cyclos		bicyclo(3.2.0)hep-tane is part
			of a polycyclo ring system
			having at least three cyclos

306	Plural 1-thia-4-aza- bicyclo(3.2.0)hep-tane ring	323	Base salt formation of 3-position -COOH group
	systems attached directly or indirectly to each other by	324	Extracting solid from solution
307	nonionic bondingSpiro	325	The nitrogen is part of a hetero ring
308	The 6-position substituent contains phosphorus attached directly or indirectly to the bicyclo ring system by	326	Chalcogen, -C(=X)-, wherein X is chalcogen, or additional nitrogen bonded directly to the -C(=X)- group
309	nonionic bondingNitrogen containing hetero ring attached directly at the	327	Hetero ring or ring system bonded directly to the -C(=X)- group
	3-position of the bicyclo ring system	328	Nitrogen containing
310	Having -C(=X)-, wherein X is chalcogen, bonded directly		ring or ring system attached by carbon or acyclic carbon chain to the -C(=X)- group
211	at the 3-position of the bicyclo ring system	329	Polycyclo heterocyclic ring system in 6-position
311	Nitrogen or hydrogen bonded directly to the -C(=X)-group	330	substituentThe polycyclo ring
312	Nitrogen bonded directly at the 6-position of the bicyclo ring system		system is attached directly to a -C(=X)-NH- group, wherein X is chalcogen and substitution may be made for hydrogen only,
313	The 2-position substituent contains chalcogen, nitrogen or halogen		which group is between the polycyclo ring system and the 1-thia-4-aza
314	Having -C(=X)-, wherein X is chalcogen, single bonded directly to the nitrogen (e.g., penicillin F, etc.)	331	<pre>bicyclo(3.2.0)hep-taneAcyclic nitrogen or azide attached indirectly to the -C(=X)- group by acyclic</pre>
315	Processes utilizing penam containing compound	332	nonionic bonding
316	Introduction of - C(=X)- group, wherein X is	332	wherein X is chalcogen, bonded directly to the nitrogen
	<pre>chalcogen, onto nitrogen (e.g., carboxamide formation,</pre>	333	Hetero ring bonded directly to the -C(=X)- group
317	etc.)Boron, silicon or	334	
318	phosphorus containing reactantEsterification of the		bonded directly to the $-C(=X)$ -group
	3-position -C(=X)X- group, wherein the X's may be the same or diverse chalcogens	335	Additional acyclic nitrogen or acyclic chalcogen in the 6-position substituent
319	Sulfur-oxidation, epimerization, 6-alkoxylation, de-esterification or reduction	336	The -C(=X)- group, an unsubstituted benzene ring and -NHH bonded directly to the
320	or anhydrous forms, or special	225	<pre>same carbon atom (e.g., ampicillin, etc.)</pre>
321	crystalline formsConversion of amine	337	Cycloaliphatic ring in 6-position substituent
322	salts to metal salts	338	Benzene or hetero ring in 6-position substituent
344	Purification utilizing solid adsorbent	339	

340		357	<pre>Having -C(=X)-, wherein X is chalcogen, bonded directly to the additional chalcogen</pre>
241	chain between the ring and the $-C(=X)$ - group	358	The additional chalcogen is sulfur which is bonded
341		359	<pre>directly to chalcogenThe sulfur is double bonded directly to the chalcogen</pre>
342	Unsubstituted hydrocarbyl chain between the ring and the -C(=X)- group	360	Additional carbon bonded directly to the additional chalcogen
343	Amine addition salts of 3-position -COOH group	361	Halogen attached directly at the 4-position of the hetero
344	Nitrogen containing		ring by nonionic bonding
	hetero ring in the cation (i.e., amine moiety)	362	The 4-position of the hetero ring is unsubstituted or alkyl
345	Plural nitrogens in		substituted only
346	<pre>the cation (i.e., amine moiety)Processes</pre>	363	Nitrogen bonded directly at the 3-position of the hetero
347		264	ring
347	<pre>Bicyclo ring system which is 1-oxa-4-aza- bicyclo(3.2.0)hep-tane</pre>	364	Nitrogen bonded directly at the 3-position of the hetero ring
	(including unsaturated)	450	The hetero ring contains at
348	Acyclic carbon double	430	_
340			least eight members including
	bonded directly at the 2-	4.5.1	nitrogen and carbon
	position of the bicyclo ring	451	Chalcogen double bonded
2.40	system		directly to a ring carbon of
349	Chalcogen attached		the hetero ring which is
	directly by a single bond to		adjacent to the ring nitrogen
	the carbon or to an acyclic		(e.g., laurolactam, etc.)
	carbon chain which contains	452	Heavy metal, aluminum, boron
	the carbon		or silicon containing
350	The ring system is 4-aza-	453	Spiro
	bicyclo(3.2.0)heptane	454	Chalcogen in the hetero ring
	(including unsaturated) and	455	Polycyclo ring system which
	has sulfur bonded directly at		contains the hetero ring as
	the 2-position		one of the cyclos
351	Thienamycin per se or salt	456	Two of the cyclos share at
	thereof		least three ring members or a
352	Five-membered hetero ring		ring member is shared by three
	consisting of one nitrogen,		of the cyclos (e.g., bridged,
	one sulfur and three carbons		peri-fused, etc.)
	as one of the cyclos of the	457	A five-membered cyclo of
	polycyclo ring system		the polycyclo ring system
353	Double bond between ring		consists of four ring carbons
	members of the five-membered		and one ring oxygen (e.g.,
	hetero ring		fused rifamycins, etc.)
354	Additional chalcogen bonded	458	Tetracyclo ring system
	directly to the hetero ring		which contains the hetero ring
355	The additional chalcogen is		as one of the cyclos (e.g.,
	bonded directly to the ring		rifamycin S, etc.)
	nitrogen	459	Nitrogen, sulfur or
356	The additional chalcogen is		halogen attached directly to
	double bonded directly to the		the tetracyclo ring system by
	hetero ring		nonionic bonding
	<u> </u>		

460	Plural nitrogens in the	478	Containing additional
4.6.1	hetero ring		heterocyclic polycyclo ring
461	Polycyclo ring system which		<pre>system having plural ring nitrogens (e.g., vinblastine,</pre>
	contains the hetero ring as one of the cyclos		vincristine, etc.)
462	Oxirane ring is one of the	479	Tricyclo ring system which
402	cyclos in the polycyclo ring	1//	contains the hetero ring as
	system (e.g., maytansinol,		one of the cyclos
	etc.)	480	Additional hetero ring
463	Nitrogen or additional	100	attached directly or
403	chalcogen attached directly to		indirectly to the hetero ring
	the hetero ring by nonionic		by nonionic bonding
	bonding	481	The additional hetero ring is
464	Utilizing oximes, oxime	101	six-membered and contains
101	salts, hydroxylamines,		nitrogen
	hydroxylamine salts or	482	Chalcogen or nitrogen attached
	nitrosating agents to form the	102	directly to the hetero ring by
	hetero ring (i.e., formation		nonionic bonding
	of the lactam ring)	483	Plural nitrogens attached
465	Heavy metal or aluminum		indirectly to the hetero ring
	containing		by acyclic nonionic bonding
466	Spiro	484	The hetero ring contains seven
467	The hetero ring contains		members including nitrogen and
	chalcogen		carbon
468	Polycyclo ring system which	485	Chalcogen double bonded
	contains the hetero ring as		directly to a ring carbon
	one of the cyclos		adjacent to the ring nitrogen
469	Plural nitrogens in the		(e.g., caprolactam, etc.)
	hetero ring	486	Heavy metal or aluminum
470	The hetero ring contains		containing
	plural nitrogens	487	Silicon or phosphorus
471	Polycyclo ring system which		attached directly or
	contains the hetero ring as		indirectly to the hetero ring
	one of the cyclos		by nonionic bonding
472	Two of the cyclos share at	488	Chalcogen in the hetero ring
	least three ring members or a	489	Plural nitrogens in the
	ring member is shared by three		hetero ring
	of the cyclos (e.g., bridged,	490	Bicyclo ring system having
	<pre>peri-fused, etc., toxiferin)</pre>		the hetero ring as one of the
473	Bicyclo ring system which		cyclos
	contains the hetero ring as	491	The chalcogen and the
	one of the cyclos		nitrogen are in the 1,5-
474	The hetero ring contains at		positions of the bicyclo ring
	least three nitrogens		system (e.g., 1,5-
475	Nitro bonded directly to		benzothiazepinone, etc.)
	ring nitrogen of the hetero	492	Plural nitrogens in the
	ring (e.g., HMX, etc.)		hetero ring
476	Polycyclo ring system which	493	Tetracyclo ring system
	contains the hetero ring as		having the hetero ring as one
	one of the cyclos	40.4	of the cyclos
477	Two of the cyclos share at	494	Nitrogen of the hetero ring
	least three ring members or a		is shared by an additional
	ring member is shared by three		cyclo of the tetracyclo ring
	of the cyclos (e.g., bridged,		system
	peri-fused, etc.)		

495	Tricyclo ring system having the hetero ring as one of the cyclos	512	<pre>Chalcogen attached indirectly to nitrogen of the hetero ring by acyclic</pre>
496	<pre>Nitrogen of the hetero ring is shared by an additional cyclo of the tricyclo ring system</pre>	513	<pre>nonionic bondingSulfur, -C(=X)-, wherein X is chalcogen, or nitrogen, other than as nitro or</pre>
497	<pre>Additional hetero atom in the additional cyclo of the tricyclo ring system</pre>		<pre>nitroso, bonded directly to the carbocyclic ring of the bicyclo ring system</pre>
498	The additional cyclo is five-membered consisting of nitrogen and carbon (e.g.,	514	Nitrogen in the 1-position substituent of the bicyclo ring system
499	<pre>imidazobenzodiazepinones, etc.)The additional cyclo</pre>	515	Preparation by cyclizing benzophenones or imine derivatives thereof
499	consists of three nitrogens and two carbons (e.g., triazolobenzodiazepinones,	516	Preparation from a compound containing a different hetero ring
500	etc.)Bicyclo ring system having	517	The bicyclo ring system is 1,5-benzodiazepine (including
	the hetero ring as one of the cyclos	518	hydrogenated)Additional chalcogen
501	At least three nitrogens in the hetero ring		double bonded directly to ring carbon of the hetero ring
502	At least three hetero atoms in the bicyclo ring system	519	Polycyclo ring system which contains the hetero ring as
503	Chalcogen in the bicyclo ring system	520	one of the cyclosTwo of the cyclos share at
504	The bicyclo ring system is 1,4-benzodiazepine (including hydrogenated)		least three ring members or a ring member is shared by three of the cyclos (e.g., bridged,
505	The chalcogen double bonded directly to the hetero ring is sulfur	521	<pre>peri-fused, etc.)Plural hetero atoms in the polycyclo ring system</pre>
506	Additional chalcogen bonded directly to ring carbon of the hetero ring	522	Tricyclo ring system which contains the hetero ring as one of the cyclos
507	The additional chalcogen is bonded directly at the 3-position of the bicyclo ring	523	Bicyclo ring system which contains the hetero ring as one of the cyclos
508	<pre>systemNitrogen or -C(=X)-,</pre>	524	Additional hetero ring containing
	wherein X is chalcogen, attached indirectly to the	525	Plural seven-membered hetero rings
	chalcogen by acyclic nonionic bonding	526	Additional chalcogen bonded directly to the hetero ring
509	Acyclic nitrogen bonded directly to the hetero ring	527	Nitrogen bonded directly to the hetero ring
510	Having $-C(=X)-$, wherein X is chalcogen, bonded directly	528	The nitrogen is bonded additionally only to hydrogen
511	<pre>to the hetero ringHalogen bonded directly to the hetero ring</pre>	529	Having -C(=X)-, wherein X is chalcogen, bonded directly to the hetero ring
		530	Halogen bonded directly to the hetero ring

531	Chalcogen or nitrogen attached indirectly to the	551	Nitrogen bonded directly to ring carbon of the hetero ring
	hetero ring by nonionic bonding	552	Bicyclo ring system which contains the hetero ring as
532	Preparing from a compound containing a hetero ring	553	one of the cyclosThe hetero ring contains
533	The hetero ring is a lactam (i.e., $-C(=X)-NH-$ is part of		plural nitrogens (e.g., 1,3-diazepines, etc.)
	the ring, wherein X is chalcogen and substitution may	554	The hetero ring contains at least three nitrogens
534	be made for the hydrogen only)Preparing from a compound containing a cycloaliphatic	555	Polycyclo ring system which contains the hetero ring as one of the cyclos
	ring	556	Two of the cyclos share at
535	<pre>The reactant is a cyclic oxime</pre>		least three ring members or a ring member is shared by three
536 537	Gas phase rearrangementAcyclic -C(=X)X-, wherein		of the cyclos (e.g., bridged, peri-fused, etc.)
	the X's are the same or diverse chalcogens, attached directly to the cycloaliphatic	557	<pre>Tricyclo ring system which contains the hetero ring as one of the cyclos</pre>
538	ring by nonionic bondingCyclization to form the	558	Nitrogen of the hetero ring is shared by an additional
539	hetero ringReactant contains a cyano		cyclo of the tricyclo ring system
	group	559	The additional cyclo has
540	Purification or recovery		at least six ring members
541	Heavy metal or boron containing	560	Chalcogen in the tricyclo ring system
542	Phosphorus attached directly or indirectly to the hetero ring by nonionic bonding	561	The additional cyclo consists of one nitrogen and four carbons (e.g.,
543 544	Spiro	562	diazepinoindoles, etc.)The additional cyclo
545	The hetero ring contains chalcogenPlural nitrogens in the	302	consists of two nitrogens and three carbons (e.g.,
313	heterocyclic ring		imidazobenzodiazepines, etc.)
546	Polycyclo ring system which contains the hetero ring as one of the cyclos	563	<pre>s-Triazolo(4,3-a)(1,4)- benzodi-azepines (including hydrogenated)</pre>
547	Tricyclo ring system which contains the hetero ring as one of the cyclos	564	Chalcogen, nitrogen, cyano or halogen bonded directly to ring carbon of the
548	At least three ring hetero atoms in the tricyclo ring system	565	<pre>triazolo ringNitrogen attached indirectly to ring carbon of</pre>
549	Sulfur and nitrogen are bonded directly to each other	566	the triazolo ring by acyclic nonionic bondingThe unshared ring carbon
550	<pre>in the hetero ringThe nitrogen of the hetero ring is bonded directly to</pre>	500	of the triazolo ring is unsubstituted or alkyl
	both remaining rings of the tricyclo ring system (e.g., dibenzo(b,e)(1,4)thiazepine, etc.)	567	<pre>substituted onlyBicyclo ring system which contains the hetero ring as one of the cyclos</pre>

568	At least three ring hetero atoms in the bicyclo ring system	585	Chalcogen or nitrogen attached indirectly to ring nitrogen of the bicyclo ring
569	1,4-benzodiazepines (including hydrogenated)		system by acyclic nonionic bonding
570	<pre>Chalcogen bonded directly to ring carbon of the hetero ring</pre>	586	<pre>Tricyclo ring system which contains the hetero ring as one of the cyclos</pre>
571	Nitrogen bonded directly to ring carbon of the hetero ring	587	The hetero ring shares ring members with each of two benzene rings in the tricyclo
572	Chalcogen or nitrogen attached indirectly to ring carbon of the hetero ring by	588	<pre>ring system (e.g., morphanthridines, etc.)The nitrogen of the hetero</pre>
573	acyclic nonionic bondingChalcogen or nitrogen attached indirectly to ring		ring is bonded directly to each of the two benzene rings (e.g., iminodibenzyl, etc.)
	carbon of the hetero ring by acyclic nonionic bonding	589	Having -C(=X)-, wherein X is chalcogen, bonded directly
574	Formation of the 1,4- benzodiazepine ring system		to ring nitrogen of the tricyclo ring system
575	The nitrogens are in the 1,4- positions of the hetero ring	590	or indirectly to ring carbon
576	Polycyclo ring system which contains the hetero ring as one of the cyclos	591	of the hetero ring by acyclic nonionic bondingChalcogen attached
577	Plural nitrogens in the polycyclo ring system		directly or indirectly to the hetero ring by acyclic
578	<pre>Three or more hetero atoms in the polycyclo ring system</pre>	592	nonionic bondingNitrogen attached
579	<pre>Nitrogen of the hetero ring is shared by an additional cyclo of the polycyclo ring</pre>	502	indirectly to ring nitrogen of the hetero ring by acyclic nonionic bonding
580	<pre>systemThe seven-membered hetero</pre>	593	Bicyclo ring system which contains the hetero ring as
	ring shares ring members with one other cyclo only	594	<pre>one of the cyclos3-Benzazepines (including hydrogenated)</pre>
581	Two of the cyclos share at least three ring members or a ring carbon is shared by three	595	Benzene ring bonded directly to ring carbon of the hetero ring
582	<pre>of the cyclos (e.g., bridged, peri-fused, etc.)Bicyclo ring system which</pre>	596	Additional hetero ring attached directly or
	contains the hetero ring as one of the cyclos (e.g., 3-		indirectly to the hetero ring by nonionic bonding
	<pre>azabicyclo-(3.2.2)nonanes, etc.)</pre>	597	The additional hetero ring is six-membered and contains nitrogen
583	<pre>Having -C(=X)-, wherein X is chalcogen, bonded directly to ring nitrogen of the</pre>	598	Plural hetero atoms in the additional hetero ring
584	bicyclo ring systemChalcogen or nitrogen	599	Polycyclo ring system having the additional hetero
	attached directly to ring nitrogen of the bicyclo ring system by nonionic bonding	600	ring as one of the cyclosThe additional hetero ring is 1,3-diazine (including hydrogenated)

601	The additional hetero ring is 1,3-diazine (including
602	<pre>hydrogenated)The additional hetero ring is five-membered and contains nitrogen</pre>
603	Plural hetero atoms in the additional hetero ring
604	Chalcogen attached directly to the hetero ring by nonionic bonding
605	Nitrogen attached directly to the hetero ring by nonionic bonding
606	Chalcogen, additional nitrogen, or -C(=X)-, wherein X is chalcogen, attached directly to the nitrogen by nonionic bonding
607	Having -C(=X)-, wherein X is chalcogen, bonded directly to nitrogen of the hetero ring
608	Chalcogen bonded directly to the -C(=X)- group
609	Chalcogen or nitrogen attached indirectly to the hetero ring by acyclic nonionic bonding
610	The chalcogen or nitrogen is multiple bonded to a carbon (e.g., cyano or carbonyl groups, etc.)
611	Benzene ring bonded directly to the hetero ring
612	The hetero ring is unsubstituted or alkyl substituted only

FOREIGN ART COLLECTIONS

FOR 000 CLASS-RELATED FOREIGN DOCUMENTS